

# **D**ETERMINATION OF CHEMICAL COMPOSITION AND SCAVENGING EFFECTIVENESS OF LEAF EXTRACTS OF *TERMINALIA CATAPPA* IN GEIDAM LOCAL GOVERNMENT AREA OF YOBE STATE, NIGERIA.

**MR. MOHAMMED AUDU, MR. MUSA MOHAMMED GARANDAWA AND MR. UMAR DAN AZUMI**

*School of Science, Department of science laboratory technology; Mai Idris Aloomo Polytechnic P.M.B 1020 Geidam, Yobe state, Nigeria.*

## **ABSTRACT**

*Medicinal plants have been identified and used throughout human history, plants have ability to synthesize a wide variety of chemical compounds. Many of which are efficacious and contain compounds that are potential drugs which require further examinations. Plants are rich in a wide variety of secondary metabolites such as tannins, terpenoids, alkaloids and flavonoids etc., which have been found to possess to antimicrobial properties in vitro. Antioxidants are molecules that inhibit or quench free radical reactions and delay or inhibit cellular damage. Although, almost all organisms possess antioxidant defence and repair systems that have evolved to protect them against oxidative damage, these systems are insufficient to prevent the*

## **Introduction:**

Medicinal plants have been in used throughout human history to treat aliment and diseases. Plants have the ability to synthesize a wide variety of chemical compounds (Lai and Roy, 2004). Many of which are efficacious and contain substances that are potential drugs that require further examinations (Abdulrahman et al., 2007). Plants have evolved with the ability to synthesize chemical compounds that can help them defend against attack from a wide variety of predators such as

damage entirely. However, antioxidant supplements, or foods containing antioxidants, may be used to help human body reduced oxidative damage. Studies on the nutritional value and biological activity of the kernel of *Terminalia catappa* revealed that it has a good digestibility, exerts a strong antioxidant activity, possesses anti-HIV properties, anti-asthma properties, anti-inflammatory, anti-carcinogenic, antibacterial and hepatoprotective properties. The fresh leaf of *Terminalia catappa* was air dried under laboratory condition, was ground into fine powder extracted with methanol using cold infusion method and partitioned using solvent of gradient polarities such as *n*-hexane, ethyl acetate and *n*-butanol. Methanol crude extract, *n*-hexane portion, ethyl acetate portion, *n*-butanol portion and aqueous portion revealed the presence of carbohydrate, cardiac glycoside, flavonoid, terpenoids, saponins, tannins and alkaloid. However, soluble starch, phlobatannins and glycosides were not found in the extracts. The methanol crude extract showed the percentage inhibition of 98.25 at 10 $\mu$ g/ml, 97.40 at 20 $\mu$ g/ml, 96.94 at 30 $\mu$ g/ml, 96.63 at 40 $\mu$ g/ml and 97.10 at 50 $\mu$ g/ml; *n*-butanol portion showed the percentage inhibition of 95.75 at 10 $\mu$ g/ml, 96.40 at 20 $\mu$ g/ml, 96.15 at 30 $\mu$ g/ml, 96.40 at 40 $\mu$ g/ml and 96.15 at 50 $\mu$ g/ml; *n*-hexane portion showed the percentage inhibition of 95.50 at 10 $\mu$ g/ml, 95.65 at 20 $\mu$ g/ml, 95.80 at 30 $\mu$ g/ml, 95.75 at 40 $\mu$ g/ml and 95.75 at 50 $\mu$ g/ml; ethyl acetate portion showed the percentage inhibition of 78.35 at 10 $\mu$ g/ml, 87.65 at 20 $\mu$ g/ml, 95.00 at 30 $\mu$ g/ml, 94.75 at 40 $\mu$ g/ml and 94.70 at 50 $\mu$ g/ml and the aqueous portion showed the percentage inhibition of 94.40 at 10 $\mu$ g/ml, 95.10 at 20 $\mu$ g/ml, 96.00 at 30 $\mu$ g/ml, 95.50 at 40 $\mu$ g/ml and 96.05 at 50 $\mu$ g/ml.

**Keywords:** *Terminalia catappa*, Phytochemicals, Proximate analysis, Elemental analysis, Secondary metabolites

Insects, fungi and herbivorous mammals (Neelavathi et al., 2013). Plants are rich in a wide variety of secondary metabolites such as tannins, terpenoids, alkaloids and flavonoids etc, which have been found in vitro to have antimicrobial properties (Dahanukar, 2000). *Terminalia Catappa* is

a large, deciduous tree with smooth grey bark and whorled branches that form a canopy and is found in tropical and subtropical regions, it is widely planted throughout the tropics as an ornamental tree for shade for the edible nuts. Terminalia catappa contains hydrolyzable tannins punicalagin (major tannin), punicalin, terflavins A and B, tergalagin, tercatanin, chebulagic acid, geraniin, granatin B, corilagin), flavonoids (isovitexin, vitexin, isoorientin, rutin) and triterpenoids (ursolic acid, 2 $\alpha$ , 3 $\beta$ , 23-trihydroxyurs-12-en-28 oic acid and asiatic acid) (Liu et al., 1996 and Kinoshita et al., 2007). The leaves, bark and fruit of the tree Terminalia catappa L. (Combretaceae) have been commonly used as a folk medicine for antidiarrhea, antipyretic and haemostatic purposes (Lin, 1992). The leaves of T. catappa have been used for the prevention and treatment of hepatitis and liver-related diseases (Lin and Khan, 1990). Antioxidants are molecules that inhibit or quench free radical reactions and delay or inhibit cellular damage (Young et al., 2001). Although almost all organisms possess antioxidant defence and repair systems that have evolved to protect them against oxidative damage, these systems are insufficient to entirely prevent the damage (Simic, 1988). The aim of this study is to find out chemical composition and scavenging effectiveness of the leaf of Terminalia catappa.

## Methodology

### Sample collection and Preparation

Fresh leaf of Terminalia catappa was collected from Geidam local government of Yobe State, Nigeria, and identified by a Plant Taxonomist. The plant leaf material was air-dried in the laboratory at room temperature. The leaf of the plant was ground to fine powder using wooden mortar and pestle and the sample was given a voucher number (TT99), stored in the laboratory Department of science laboratory technology (S.L.T) Mai Idris Aloomo Polytechnic Geidam for further analysis.

### Sample extraction and Partitioning

The ground leaf material (800g) was extracted with 85% methanol using maceration (cold infusion) method for 72 hours. The crude extract was

concentrated under reduced temperature. The crude extract was then stored in a desiccator. The methanol extract of *Terminalia catappa* was partitioned with n-hexane until exhaustion. The aqueous fraction was partitioned with ethyl acetate and also partitioned exhaustively with n-butanol. The n-butanol portion and the aqueous portion were then evaporated using rotatory oven.

### Phytochemical Screening of *Terminalia catappa*

Phytochemical screening was carried out on the methanol crude extract and the partitioned portions of the plant *Terminalia catappa* using standard protocols as described by (Trease and Evans, 2002, Silver et al., 1998, Markham, 1987, Brain and Turner, 1975).

### Antioxidant Studies

The free radical scavenging capacity of the extracts was determined using 1,1-Diphenyl 1,2- Picrylhydrazyl (DPPH) (Li et al., 2001).

### Results and Discussion

**Table 1: Phytochemical Screening of *Terminalia catappa* Methanol Crude extract, n-Hexane Portion, Ethyl acetate Portion, n-Butanol Portion, and Aqueous Portion.**

S/NO.	Test	MCE	NHP	EAP	NBP	AQP
1.	Carbohydrate	+	-	-	+	+
2.	Soluble starch	-	-	-	-	-
3.	Phlobotannins	-	-	-	-	-
4.	Glycosides	-	-	-	-	-
5.	Cardiac glycoside	+	-	-	+	+
6.	Flavonoid	+	-	-	+	+
7.	Terpenoid	+	+	+	-	+
8.	Saponins	+	-	-	-	+
9.	Alkaloid	+	-	-	+	-
10.	Tannins	+	-	-	+	+

**Key:** MCE = Methanol Crude Extract, NHP = n-Hexane Portion, EAP = Ethyl Acetate Portion, NBP= n-Butanol Portion, AQP = Aqueous Portion, (+) = Present and (-) = Absent.

Methanol crude extract, n-hexane portion, ethyl acetate portion, n-butanol portion and aqueous portion revealed the presence of carbohydrate, cardiac glycoside, flavonoid, terpenoids, saponins, tannins and alkaloid. However, soluble starch, phlabotannins and glycosides were not found in the extracts. The presence of secondary metabolites in the Terminalia catappa such as tannins, cardiac glycoside, flavonoid, saponins and Phenolic compound, indicates or implicate the medicinal value of it. These compounds have been reported to have antioxidants property and exhibit a wide range spectrum of medicinal properties such as anti-cancer, anti-inflammatory and anti-diabetes (Odukoya et al., 2005, Abdulrahman et al., 2010).

**Table 2: Percentage Inhibition of Standard, Methanol extract, n-Butanol Portion, n-Hexane Portion, Ethyl acetate Portion, and Aqueous Portion of Terminalia catappa**

S/NO	Extract	10 µg/ml	20 µg/ml	30 µg/ml	40 µg/ml	50 µg/ml
1	VIT C	99.25±0.05000 <sup>a</sup>	98.95±0.05000 <sup>a</sup>	98.85±0.1500 <sup>a</sup>	96.65±1.850 <sup>a</sup>	98.25±0.250 <sup>a</sup>
2	ME	98.25±0.1500 <sup>a</sup>	97.40±0.2000 <sup>a</sup>	96.94±0.03500 <sup>a</sup>	96.63±0.035 <sup>a</sup>	97.10±0.1000 <sup>a</sup>
3	NBP	95.75±0.3500 <sup>a</sup>	96.40±0.3000 <sup>a</sup>	96.15±0.7500 <sup>a</sup>	96.40±1.000 <sup>a</sup>	96.15±1.050 <sup>a</sup>
4	NHP	95.50±0.9000 <sup>a</sup>	95.65±1.550 <sup>a</sup>	95.80±1.700 <sup>a</sup>	95.75±1.950 <sup>a</sup>	95.75±2.150 <sup>a</sup>
5	EAP	78.35±0.95 <sup>b</sup>	87.65±0.7500 <sup>a</sup>	95.00±1.200 <sup>a</sup>	94.75±1.450 <sup>a</sup>	94.70±0.900 <sup>a</sup>
6	AQP	94.40±0.0 <sup>a</sup>	95.10±0.3000 <sup>a</sup>	96.00±0.1000 <sup>a</sup>	95.50±0.1000 <sup>a</sup>	96.05±0.150 <sup>a</sup>

**Key:** VIT C = Vitamin C (Ascorbic acid), ME = Methanol extract, NBP = n-Butanol Portion, NHP = n-Hexane Portion, EAP = Ethyl Acetate Portion, and AQP = Aqueous Portion.

a = P ≤ 0.05 significance across the column

b = P ≤ 0.05 insignificance across the column

## Conclusions

The results of antioxidant evaluation showed that the methanol crude extract as well as the partitioned portions indicates promising antioxidant properties of leaf of Terminalia catappa, the reducing power of the extracts compared with the reference standard antioxidant drug (ascorbic acid). The DPPH antioxidant assay is based on the ability of DPPH, a stable free radical, to decolorize in the presence of antioxidants. This study has already shown that the leaf of Terminalia catappa contains many medicinal and important phytochemicals.

## Reference

- Abdullahi Mann, Kolo Ibrahim, Adebayo O. Oyewale, Joseph O. Amupitan, Majekodumi O. Fatope and Joseph I. Okogun(2012). Isolation and Elucidation of Three Triterpenoids and Its Antimycobacterial Activity of Terminalia Avicennioides. *Am. J. of Organic Chemistry*, 2(2): 14-20.
- Aimola, I. A., Inuwa, H. M., Mamman, A. I., Habila, N., Agbaji, A. S. and Omoniwa, D. (2011). Terminalia catappa extract enhances erythropoiesis in adult balb C mice. *J. of molecular boil. Research*.1(1):40-46.
- Abdulrahman, F.I., Ogugbuaja, V.O., Onyeyili, P.A. (2007). Phytochemical screening and elemental content of the root bark extract of Vitex doniana Sweet. *Bulletin of Pure and App. Sci.* 26C (1):55.
- Aneja, K. R., Sharma, C., and Joshi, R. (2012). Antimicrobial activity of Terminalia arjuna Wight and Arn: An ethno medicinal plant against pathogens causing ear infection. *Brazilian J. of otorhinolaryngology*. 78(1):68-74.
- Atnasooriya, W. D. and Dharmasiri, M. G. (2000). Effect of Terminalia catappa on sexual behaviour and fertility of male rats. *Asian J. of Andrology*, 2, 213-219.
- Babayi, H., Kolo, I., Okogun, J. I. and Ijah, U. J. J. (2004). The antimicrobial activities of methanolic extracts of Eucalyptus camaldulensis and Terminalia catappa against some pathogenic microorganisms. *Biokemistri*, 16(2):106-111.
- Bag, A., Bhattacharya, S. K., and Chattopadhyay, R. R. (2013). The development of Terminalia chebula Retz.(Combretaceae) in clinical research. *Asian Pac. J. Tropical Biomed.*, 3(3):244- 252.
- Bayani, U., Singh A. V., Zamboni, P., and Mahajan, R. T. (2009). Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options *Current neuropharmacology*, 7(1): 65-74.
- Bennicelli, R., Stepniewska, Z., Banach, A., Szajnocha, K. and Ostrowski, J. (2004). The ability of Azolla caroliniana to remove heavy metals Hg(II), Cr(III), Cr(VI) from municipal waste water. *Chemosphere*. 55:141-146.
- Bulama, J. S., Dangoggo, S. M., Halilu M. E., Tsafe A. I. And Hassan S. W. (2014). Isolation and Characterization of Palmitic Acid from Ethyl Acetate Extract of Root Bark of Terminalia Glaucescens. *Chemistry and Materials Resear.*, 6(12): 140-144.

- Bulama, J. S., Dangoggo, S. M. and Mathias, S. N. (2015). Isolation and Characterization of Beta-Sitosterol from ethyl acetate extract of root bark of *Terminalia glaucescens*. *Inter. J. of Scientific and Research Pub.*, 5(3):2250-3153.
- Carrubba, A. and Scalenghe, R. (2012). Scent of Mare Nostrum – Medicinal and Aromatic Plants (MAPs) in Mediterranean soils. *J. of the Sci. of Food and Agric.* 92 (6): 1150–1170.
- Chanda, S., Dudhtra S. and Kaneria M. (2010). Antioxidative and Antimicrobial Effects of Seeds and Fruit Rind of Nutraceutical Plants Belonging to the Fabaceae Family. *Food and Func.*, Vol.1, pp. 308-315.
- Chatha, S. A. S., Hussain, A. L., Asad, R., Majeed, M. and Aslam, N. (2014). Bioactive Components and antioxidant properties of *Terminalia arjuna* L. extracts, 5(2): 1-5.
- Chaudhari, G. M. and Mahajan, R. T. (2015). Comparative Antioxidant Activity of Twenty Traditional Indian Medicinal Plants and its Correlation with Total Flavonoid and Phenolic Content. *Inter. J. of Pharm. Sci. Review and Resear.* 30(1):105-111.
- Chaves, T. P., Cleido, P. S. and Veras, G. (2013). Seasonal variation in the production of secondary metabolites and antimicrobial activity of two plant species used in Brazilian traditional medicine. *Afri. J. of Biotech.*, 12(8): 847-853.
- Chen, P. S., Li, J. H., Liu, T. Y. and Lin, T. C. (2000). Folk medicine *Terminalia catappa* and its major tannin component, punicalagin, are effective against bleomycin-induced genotoxicity in Chinese hamster ovary cells. *Cancer Letters* 152, 115–122.
- Chu, S. C., Yang, S. F., Liu, S. J., Kuo, W. H., Chang, Y. Z. and Hsieh, Y. S. (2007). In vitro and in vivo antimetastatic effects of *Terminalia catappa* L. leaves on lung cancer cells. *Food and chemical tox.*, 45(7), 1194-1201.
- Sunita Bhatnagar and Rituparna Panigrahi, (2015). Cytotoxic, phytochemical and antioxidant potential of marshy herb *Enhydra fluctuans* Lour. *International Journal of Ethnobiology & Ethnomedicine.* 1(1), 1-6
- Sun, H. D., Huang, S. X. and Han, Q. B. (2006). Diterpenoids from *Isodon* species and their biological activities. *Nat. Prod. Reports*, 23, 673-698.
- Subramanian, R., Subbramaniyan, P., and Raj, V. (2014). Phytochemical Screening, Total Phenolic Contents and Antioxidant Activity of *Syzygium caryophyllatum* and *Syzygium densiflorum*, *J. of Biologically Active Products from Nat.*, 4(3):224-235.
- Sodipo, O. A., Abdulrahman, F. I., Akan, J. C., and Akinniyi, J. A. (2008). Phytochemical screening and elemental constituents of the fruit of *Solanum macrocopum* Linn. *Continental J. Appl. Sci.*, 3: 88-97.
- Siaka Sohro, Kra Adou Koffi Mathieu, Ahon Gnamien Marcel, kassi Amian Brise Benjamin, Ouattara Sitapha, Aw Sadat, Coulibaly Adama, Soro Yaya and Djaman Allico Joseph (2015). Antifungal Activity of *Terminalia superba* (Combretaceae). *Journal of Experimental Biology and Agricultural Sciences.* 3(2), 163-173
- Savithramma, N., Linga Rao, M. and Suhrulatha, D. (2011). "Screening of Medicinal Plants for Secondary Metabolites." *Middle-East J. Sci. Res.*, 8: 579-84.
- Samir, M. El-Ameen, Laila, A. Refahy, Maher, A. Mahmoud, Amal, M. Saad, Allia, M. Abdo and Asmaa S. Mohammed. (2013). Chemical Investigation and Antioxidant Activity of Phenolic Acids from the Leaves of *Terminalia arjuna*. *Global J. of Pharm.*, 7 (4): 448-456.

- Sabu, M. C. and Kuttan, R. (2009). Antidiabetic and antioxidant activity of *Terminalia bellerica* Roxb. *Indian J. Exp. Biology*, 47: 270-275.
- Rao, B. K., Sudarshan, P. R., Rajasekhar, M. D., Nagaraju, N. and Rao, C. A. (2003). Antidiabetic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. *J. of Ethnopharmacology*; 85(1):169-172.
- Pellati, F., Benvenuti, S., Magro, L., Melegari, M. and Soragni, F. (2004). Analysis of phenolic compounds and radical scavenging activity of *Echinacea* spp. *J. Pharm. Biomed. Anal.* 35:289-301.
- Okpuzor, J., Adebessin O., Henrietta O. H. and Amadi I. (2008). The Potential of medicinal plants in sickle cell disease control: A review. *Inter. J. of Biomed.and Health Sci.*, 4(2); 47-55.
- Oduro, I., Larbie, C., Amoako, T. N. E. and Antwi-Boasiako A. F. (2009). Proximate composition and basic phytochemical assessment of two common varieties of *Terminalia catappa*. *J. of Sci. and Tech.*, 29 (2): 1-7.
- Nwosu, F. O., Dosumu, O. O. and Okocha, J. O. C. (2008). The potential of *Terminalia catappa* (Almond) and *Hyphaene thebaica* (Dum palm) fruits as raw materials for livestock feed. *Afr. J. of Biotech.*, 7(24): 4576-4580.
- Nita Yadava, Rajesh Yadava and Anju Goyal (2014). Chemistry of Terpenoids. *Int. J. Pharm. Sci. Rev. Res.*, 27(2), 272-278.
- McCullough, M. L., Peterson, J. J., Patel, R., Jacques, P. F., Shah, R. and Dwyer, J. T. (2012). Flavonoid Intake and Cardiovascular Disease Mortality in a Prospective Cohort of US Adults. *The Am. J. of Clinical Nutrition*, 95 (2): 454-64.
- Desobry S. (2009). Composition and nutritional properties of seeds and oil from *Terminalia catappa* L., *Advance J. of Food Sci. and Tech.*, 1(1): 72-77.
- Masuda, T., Yonemori, S., Oyama, Y., Takeda, Y., Tanaka, T., Andoh, T., Shinohara, A. and Nakata, M. (1999). Evaluation of the antioxidant activity of environmental plants: activity of the leaf extracts from seashore plants. *J. of Agric. and Food Chemistry*, 47, 1749-1754.